

RESEARCH
REVIEW



Quantifying the conservation value of seascape connectivity: a global synthesis

Andrew D. Olds^{1,2*}, Rod M. Connolly¹, Kylie A. Pitt¹, Simon J. Pittman^{3,4}, Paul S. Maxwell^{1,5}, Chantal M. Huijbers^{1,2}, Brad R. Moore⁶, Simon Albert⁷, David Rissik⁸, Russell C. Babcock⁹ and Thomas A. Schlacher²

¹Australian Rivers Institute – Coast and Estuaries, and School of Environment, Griffith University, Gold Coast, Qld 4222, Australia, ²School of Science and Engineering, University of the Sunshine Coast, Maroochydore, Qld 4558, Australia, ³Biogeography Branch, National Oceanic and Atmospheric Administration, Silver Spring, MD 20910, USA, ⁴Marine Institute, Plymouth University, Plymouth PL4 8AA, UK, ⁵Healthy Waterways, Brisbane, Qld 4004, Australia, ⁶Coastal Fisheries Programme, Secretariat of the Pacific Community, Noumea 98848, New Caledonia, ⁷School of Civil Engineering, The University of Queensland, Brisbane, Qld 4072, Australia, ⁸National Climate Change Adaptation Research Facility, Griffith University, Gold Coast, Qld 4222, Australia, ⁹Marine and Atmospheric Research, Commonwealth Scientific and Industrial Research Organization, Perth, Qld 4102, Australia

*Correspondence: Andrew D. Olds, Australian Rivers Institute – Coast and Estuaries, and School of Environment, Griffith University, Gold Coast, Qld 4222, Australia. E-mail: aolds@usc.edu.au

ABSTRACT

Aim Connectivity structures populations, communities and ecosystems in the sea. The extent of connectivity is, therefore, predicted to also influence the outcomes of conservation initiatives, such as marine reserves. Here we review the published evidence about how important seascape connectivity (i.e. landscape connectivity in the sea) is for marine conservation outcomes.

Location Global.

Methods We analysed the global literature on the effects of seascape connectivity on reserve performance.

Results In the majority of cases, greater seascape connectivity inside reserves translates into better conservation outcomes (i.e. enhanced productivity and diversity). Research on reserve performance is, however, most often conducted separately from research on connectivity, resulting in few studies (< 5% of all studies of seascape connectivity) that have quantified how connectivity modifies reserve effects on populations, assemblages or ecosystem functioning in seascapes. Nevertheless, evidence for positive effects of connectivity on reserve performance is geographically widespread, encompassing studies in the Caribbean Sea, Florida Keys and western Pacific Ocean.

Main conclusions Given that research rarely connects the effects of connectivity and reserves, our thesis is that stronger linkages between landscape ecology and marine spatial planning are likely to improve conservation outcomes in the sea. The key science challenge is to identify the full range of ecological functions that are modulated by connectivity and the spatial scale over which these functions enhance conservation outcomes.

Keywords

Conservation planning, ecological processes, ecosystem functioning, landscape ecology, marine reserves.

INTRODUCTION

We are like islands in the sea, separate on the surface but connected in the deep.

William James

Movements of organisms, matter and energy are key ecological processes, connecting populations, habitats, food webs and ecosystems (Massol *et al.*, 2011; Hyndes *et al.*, 2014). This connectivity is also central to understanding population viability and the resilience of ecosystems to disturbance (Bernhardt & Leslie, 2013; Magris *et al.*, 2014). The fundamental importance

of these linkages has long been recognized (Odum, 1968), and in recent years connectivity has become an increasingly important consideration in spatial conservation planning, complementing the long-standing focus on species persistence, habitat quality and area (Moilanen *et al.*, 2009; Hodgson *et al.*, 2011; Kool *et al.*, 2013). Examples of an increasing integration of connectivity in conservation include: optimizing the placement of reserves and the spacing of reserve networks; identifying and protecting key corridors of animal migration; limiting the spread of invasive species; and promoting the restoration of fragmented landscapes (Rudnick *et al.*, 2012; Green *et al.*, 2014).

In the sea, as on land, the area, quality and spatial arrangement of habitats are key determinants of the distribution, movement, growth and survival of organisms (Irlandi & Crawford, 1997; Micheli & Peterson, 1999). Physical linkages between discontinuous habitats (either of the same or of different types) of the seafloor represent seascape connectivity (i.e. landscape connectivity in the sea); this connectivity can be quantified as functional connectivity (i.e. documented movement of organisms among patches or habitats) or structural connectivity (i.e. distribution of organisms or processes in relation to the spatial configuration of patches or habitats) (Grober-Dunsmore *et al.*, 2009). The importance of seascape connectivity has been well documented for post-settlement fish and crustaceans (i.e. individuals that have successfully recruited to a habitat or assemblage), which move among different habitats to spawn, forage or undertake ontogenetic habitat shifts (Pittman & McAlpine, 2003; Sheaves, 2009; Boström *et al.*, 2011). Moreover, seascape connectivity influences the spatial distribution of fish populations and fisheries catches and the outcomes of conservation initiatives (Nagelkerken *et al.*, 2012; Olds *et al.*, 2013).

Although connectivity is now a key consideration for marine conservation planning (Green *et al.*, 2014; Magris *et al.*, 2014), few studies actually provide empirical data about the conservation benefits of connectivity in marine ecosystems. There is a new urgency to address this knowledge gap because human activity has altered the condition of, and level of connectivity among, coastal populations, habitats and ecosystems (Boström *et al.*, 2011; Hyndes *et al.*, 2014; Nagelkerken *et al.*, 2015). For

example, wetlands are being increasingly fragmented by coastal development, and the physical complexity of coral reefs is being eroded at a global scale with unanticipated consequences for connectivity, ecological resilience and provisioning of ecosystem services (Unsworth & Cullen, 2010; Rogers *et al.*, 2014). Furthermore, overharvesting of mobile consumers (e.g. fish, crustaceans) and the construction of barriers (e.g. dams, weirs, levees, sea walls) to animal migration has also physically degraded key connectivity pathways, affecting food webs, assemblage composition, population viability and the condition of marine ecosystems (Fig. 1) (Valentine *et al.*, 2008; Nagelkerken *et al.*, 2015).

General concepts and paradigms in seascape ecology and connectivity have been reviewed elsewhere (e.g. Grober-Dunsmore *et al.*, 2009; Boström *et al.*, 2011; Pittman & Olds, 2015), but no synopsis exists that explicitly addresses the question of whether and how connectivity enhances the effectiveness of marine reserves. Here we review studies of seascape connectivity (i.e. connectivity among habitats in seascapes) or habitat connectivity (i.e. connectivity among patches of the same habitat), focusing on post-settlement fish and crustaceans, to assess whether greater connectivity translates into consistently better conservation outcomes for populations, assemblages or ecosystem functioning for reserves. Our review examines connectivity in seascapes that lie within reserves and span reserve boundaries. Fish and crustaceans move among habitats in seascapes, are commonly managed using reserves and are, therefore, ideal model organisms for examining the role of connectivity in con-




Figure 1 Common seascapes and impacts on connectivity in marine ecosystems (coral reefs, left; estuarine wetlands, right). We illustrate three types of connectivity: (1) largely unmodified seascapes with intact connections (a, b), (2) connections that are exploited to harvest organisms as they move regularly across seascapes (c, d), and (3) human alterations of connections illustrated by engineering structures (levees, weirs) that impede movement (e, f). Photographs by A.D.O., P.S.M., S.A. in Australia (b, e, f), Papua New Guinea (d), and Solomon Islands (a, c).

ervation. We discuss implications for spatial conservation planning and we identify important knowledge gaps to be targeted in future research.

Our focus in this review is on connectivity of organisms post-settlement in reef and wetland seascapes, but the concepts and approach that we address are relevant to other seascapes. Pre-settlement processes (i.e. larval dispersal, population connectivity) are also an important consideration in marine spatial planning (e.g. facilitating network connectivity), but the potential role of larval dispersal for marine conservation has been reviewed widely (Cowen & Sponaugle, 2009; Green *et al.*, 2014; Jones, 2015). To determine the conservation value of seascape connectivity we have therefore focused on studies that have examined these effects for adult and juvenile animals.

SEASCAPE CONNECTIVITY AND MARINE CONSERVATION

Connectivity is important for conservation, mainly because populations of mobile organisms are spatially linked and adjacent ecosystems can function as linked units forming a habitat mosaic (Massol *et al.*, 2011; Green *et al.*, 2014). Connectivity can enhance the performance of individual marine reserves (Olds *et al.*, 2014) and mechanistically links habitat patches in reserves with those in other reserves, or in adjacent fished waters (Pittman *et al.*, 2014). Connectivity influences the number of organisms exported from reserves (both adults and larvae) to replenish populations in fished waters (i.e. spillover) and links populations among reserves (e.g. Freeman *et al.*, 2009; Halpern *et al.*, 2009; Harrison *et al.*, 2012). For example, spillover of fish from reserves is likely to be greatest where reserves are surrounded by contiguous habitat that provides greater connectivity (Edgar *et al.*, 2014). This process can, however, also limit the recovery of fished populations within reserves (Babcock *et al.*, 2012). Seascape connectivity within reserves and reserve networks can also promote ecosystem resilience and facilitate access to refugia (Bernhardt & Leslie, 2013; Magris *et al.*, 2014). Better-connected populations and habitats in reserves can recover more quickly from disturbance through the arrival of individuals or propagules from other locations (i.e. recolonization effects) (Beger *et al.*, 2010; Green *et al.*, 2014) or via the ecological processes (e.g. herbivory) that mobile species provide for the functioning of ecosystems (Mumby, 2006; Olds *et al.*, 2012). Connectivity also enables organisms to access refugia in higher latitudes, or in deeper, cooler water, when moving away from the thermal impacts of climate change and, therefore, provides the mechanism for linking reserves across bioregions (Magris *et al.*, 2014).

Marine reserves provide model systems for investigating the ecological importance of connectivity for conservation. Marine reserves can have significant positive effects on the density, body size, biomass and demographic parameters of harvested fish and invertebrates (Lester *et al.*, 2009; MacNeil *et al.*, 2015), particularly when reserves prohibit fishing, are well enforced, old (> 10 years), large (> 100 km²) and isolated by deep water or sand (Edgar *et al.*, 2014). Large consumer populations inside reserves

can also affect the abundance and movement of competitors and prey species, and drive trophic cascades that structure the condition of benthic communities (Babcock *et al.*, 2010; MacNeil *et al.*, 2015). Furthermore, marine reserves can promote the transfer of carbon among ecosystems, linking ecological processes and food webs across seascapes (Langlois *et al.*, 2005; Salomon *et al.*, 2008). These effects on movement ecology and ecosystem functioning provide key mechanisms through which reserves influence connectivity (Valentine *et al.*, 2008; Olds *et al.*, 2012; Pittman *et al.*, 2014).

ECOLOGICAL STUDIES OF SEASCAPE CONNECTIVITY

To evaluate the importance of seascape connectivity in conservation we compiled a database of all peer-reviewed studies that reported effects of connectivity on marine fauna and ecological processes. The ISI Web of Knowledge database was searched using all permutations of the keywords: seascape, connectivity (connect*, link*, move*) and habitat (reef, kelp, mangrove, saltmarsh, seagrass, wetland). This database was then refined (using the keywords: MPA, no-take area, protected area, conservation area, reserve, spillover) to identify studies that examined connectivity effects in reserves, and those that directly quantified both connectivity and reserve effects. Connectivity studies focused on structural (i.e. distribution of organisms or processes) and functional (i.e. movement) measures (Grober-Dunsmore *et al.*, 2009). Reserve studies were categorized based on four common criteria for performance: 'production' (i.e. abundance, density, biomass), diversity (i.e. diversity, richness, composition), ecological processes (i.e. herbivory, predation, recruitment) and spillover (i.e. movement from inside to outside reserves) (Lester *et al.*, 2009).

We identified 213 studies that reported on connectivity, of which 143 were in coral and rocky reef seascapes and 70 in wetland systems (i.e. seagrass, mangrove and saltmarsh) (Fig. 2, Appendix). Most studies were from the Caribbean Sea (56), Australia (47), the Florida Keys (19), Tanzania (18), North America (15), the Mediterranean Sea (10), the Gulf of Mexico (8) and Indonesia (7), illustrating a concentration of research in tropical seascapes (Fig. 2, Appendix S1 in Supporting Information).

One fifth ($n = 45$) of studies that examined connectivity also addressed questions of marine reserve performance (Appendix S2). Most of these focused on structural (32) or functional (9) connectivity for fishes, whilst only four measured connectivity for crustaceans. All were conducted in reef seascapes (38 in coral reef and 7 in rocky reef systems), with most studies (35) examining the conservation benefits of seascape connectivity (i.e. linkages among different habitat types) and just a few (10) exploring the importance of habitat connectivity (i.e. linkages among patches of the same habitat type). There were no consistent differences in spatial scale between studies that examined seascape or habitat connectivity, or that focused on different types of organisms. Studies of functional connectivity were, however, generally conducted over a broader spatial scale (i.e.




Figure 2 Global distribution of coral reef, seagrass, mangrove and saltmarsh ecosystems (a); marine reserves (i.e. no-take areas), marine parks (i.e. partial no-take areas) and other conservation areas (i.e. with unknown status) (b); and connectivity studies in reef and wetland seascapes (c) (Appendix). Information on the extent and geographic distribution of ecosystem types and marine conservation areas has been sourced from the United Nations Environment Programme World Conservation Monitoring Centre (UNEP-WCMC).

thousands to tens of thousands of metres) than studies of structural connectivity (i.e. hundreds to thousands of metres) (Appendix S2).

Twenty-seven studies examined connectivity effects inside reserves without explicitly testing reserve effectiveness ('connectivity effects inside reserves'). Eight studies quantified connec-

tivity and reserve effects separately (i.e. they evaluated connectivity but did not examine whether it influenced reserve performance). Ten studies (i.e. fewer than 5% of seascape studies that examined connectivity) directly evaluated whether connectivity influenced reserve performance (Fig. 3, Appendix S2). These findings show that research on marine reserve perfor-




Figure 3 Connectivity effects contrasted between study types that incorporated the role of marine reserves to varying degrees: (a) studies that examined connectivity inside reserves but did not address reserve effects per se; (b) studies that quantified connectivity and reserve effects separately; and (c) studies that tested for an interaction between connectivity and reserve effects. Circle quarters represent summaries against reserve performance measures, with the number of studies of each type shown outside (in parentheses) and the total number of studies inside. The proportion of studies reporting significant effects is illustrated by a quadrant's size (and provided as percentages) (see Appendix S2). Panel (d) illustrates some of the principal attributes and mechanisms examined, such as the size and biomass of fish populations, the composition and diversity of assemblages and trophic interactions (e.g. herbivory). Photographs by A.D.O.

mance and seascape connectivity is typically conducted separately. This may suggest a separation (in terms of ecological theory or study design) between the disciplines of conservation biology and landscape ecology in many marine ecosystems. The 27 studies that examined connectivity inside reserves reported positive effects of greater connectivity on metrics that are commonly used to assess reserve performance, including: production (94% of studies), diversity (92%), ecological processes (100%) and spillover (71%) (Fig. 3, Appendix S2). The same was true for the eight studies that quantified connectivity and marine reserve effects separately, with 75–100% of these reporting positive effects of connectivity on the reserve performance measures production, diversity and spillover (Fig. 3, Appendix S2).

The most compelling result, however, is provided by studies that were designed specifically to test for interactive effects of seascape connectivity on reserve performance. Whilst comparatively under-represented with 10 out of 45 studies, 80–100% of these reported positive effects of greater connectivity on production (i.e. density or biomass), diversity (i.e. richness or composition) or ecological processes (i.e. herbivory, predation or recruitment) in reserves, compared with fished locations and reserves supporting isolated habitats (Fig. 3, Appendix S2). In the Caribbean Sea, for example, connectivity (quantified as habitat isolation and density at the scale of 200–1000 m) influenced marine reserve effects on fish biomass, species richness and assemblage composition on coral patch reefs at Glover's Atoll, Belize (Huntington *et al.*, 2010). In the Florida Keys, con-

nectivity (quantified as habitat isolation at the scale of 0–100 m) also influenced marine reserve effects on fish abundance and assemblage composition, as well as herbivory and predation rates, in seagrass meadows adjacent to coral reefs (Valentine *et al.*, 2008). Similarly, in the western Pacific Ocean connectivity (quantified as habitat isolation at the scale of 100–1000 m) influenced marine reserve effects on fish abundance, species richness and assemblage composition on coral reefs, and in adjacent mangroves and seagrass (Olds *et al.*, 2013). These seascape connectivity effects were largely driven by the distribution and behaviour of fish species that regularly use multiple habitats (or habitat patches) through their lives (e.g. fish from the families Haemulidae, Lethrinidae, Lutjanidae, Scaridae, Siganidae). Furthermore, the synergistic impact of reserves and connectivity on herbivorous fish (family Siganidae) also enhanced herbivory and coral recruitment on inshore coral reefs in Moreton Bay, eastern Australia (Olds *et al.*, 2012). The results of studies that examined connectivity and reserve effects underscore the importance of cross-disciplinary integration of concepts and techniques to address new applied research questions at spatial scales that are operationally meaningful to conservation. We acknowledge that publication bias against neutral or insignificant results and our choice of search terms may have influenced the pool of literature reviewed (Babcock *et al.*, 2010; Magris *et al.*, 2014). These factors would, however, not detract from one of our main findings, namely that studies of reserve performance and seascape connectivity are often conducted separately.

INTEGRATING CONNECTIVITY INTO MARINE CONSERVATION

Based on the evidence available that strongly suggests a positive effect of greater connectivity on reserve performance, we recommend that seascape connectivity should be an important criterion in marine conservation. Four conditions will often have to be met when seeking to more comprehensively integrate connectivity in marine reserve design: (1) evaluate whether the benefits of connectivity result in a trade-off with other conservation considerations; (2) determine the spatial scales over which connectivity is likely to benefit reserve performance; (3) decide on a suitable approach for integrating connectivity into conservation planning and explicitly consider the effects of reserves on connectivity; and (4) quantify whether positive effects of connectivity for particular species (or groups of species) result in conservation benefits for the functioning of ecosystems, and vice versa.

Connectivity can have undesirable effects if it promotes the spread of invasive species, pathogens or pollutants (Rudnick *et al.*, 2012; Kool *et al.*, 2013). A stronger and more consistent emphasis on connectivity in conservation planning may also conflict with other socio-ecological considerations, including the representation of biodiversity, habitat area or quality (Beger *et al.*, 2010; Hodgson *et al.*, 2011), and the spatial spreading of risk to maximize population persistence and ecosystem resilience (Bernhardt & Leslie, 2013; Magris *et al.*, 2014). To be a clear priority for conservation, the potential benefits of connectivity must, therefore, exceed these risks and any potential costs to other conservation objectives.

Spatial scale is a key consideration for both marine conservation planning and seascape ecology (Magris *et al.*, 2014; Martin *et al.*, 2015). The synergistic effects of connectivity and reserves have typically (in six out of ten studies) been reported at the scale of hundreds to thousands of metres, which corresponds to tidal, daily and some ontogenetic fish movements (Boström *et al.*, 2011; Olds *et al.*, 2013), and is smaller than the size of most reserves (Lester *et al.*, 2009; Huijbers *et al.*, 2014). Within this range, the positive effects of connectivity will likely decline with distance, but the scale and shape of such effects may vary with the spatial extent of seascapes, the particular habitats these support and the mobility of the species of interest (*sensu* Olds *et al.*, 2013). The scale of any such decline-by-distance effect would also likely vary with the type of connectivity in question, with population connectivity (i.e. through larval dispersal and genetic exchange) occurring among habitat patches that are separated by much greater distances (i.e. tens to hundreds of kilometres) (e.g. Harrison *et al.*, 2012) than examined here. Studies are needed to identify factors that determine the scale over which connectivity effects reserve performance (priority questions 1 and 2, Table 1).

The identification of seascapes with optimal connectivity for maintaining biodiversity, productivity and ecosystem functioning is now a major goal in applied marine ecology (Pittman *et al.*, 2014; Nagelkerken *et al.*, 2015). With quantitative data on the scale of connectivity effects in seascapes, and the mode by

Table 1 Priority questions for research on connectivity and conservation in marine ecosystems. Cited studies provide examples of potential approaches for examining each priority question.

| Priority research questions |
|---|
| 1. In situations where positive effects of connectivity on reserve performance are evident, what is the shape of the 'distance–decline' response curve and are thresholds evident in the effect of connectivity (e.g. Martin <i>et al.</i> , 2015)? |
| 2. What factors are most important in determining the scale of connectivity effects on reserve performance (e.g. species biology, seascape composition, habitat quality, fishing pressure) (e.g. Olds <i>et al.</i> , 2013)? |
| 3. Does connectivity improve reserve performance in seascapes that do not include coral reefs (e.g. estuaries, rocky reefs, sandy beaches, seamounts) (e.g. Langlois <i>et al.</i> , 2005)? |
| 4. Does connectivity improve reserve performance for organisms other than fishes (e.g. MacArthur <i>et al.</i> , 2011)? |
| 5. How widespread and prominent are connectivity effects on ecological processes (e.g. herbivory, predation, recruitment) in reserves (e.g. Valentine <i>et al.</i> , 2008)? |
| 6. Do the effects of connectivity on ecological functions improve the resistance of protected ecosystems to disturbance, or influence their recovery (e.g. Olds <i>et al.</i> , 2012)? |
| 7. To what extent do the characteristics of seascapes at reserve boundaries influence movement within reserves and spillover from reserves (e.g. Pittman <i>et al.</i> , 2014)? |

which these affect reserve performance, seascape connectivity can be integrated into conservation by modifying reserve selection algorithms to incorporate connectivity (Moilanen *et al.*, 2009). Decision support tools (e.g. Marxan or Zonation) already provide some options for incorporating connectivity into conservation, and can be refined through further integration of quantitative data on connectivity (Beger *et al.*, 2010; Magris *et al.*, 2014). Given the potential for high spatial variability in connectivity effects, this may be achieved by prioritizing the conservation of seascapes that contain large, high-quality patches of multiple habitats (e.g. coral reefs and mangroves) within the spatial extent of species dispersal capabilities (i.e. home ranges, ontogenetic habitat shifts, pelagic larval duration) (e.g. Beger *et al.*, 2010; Edwards *et al.*, 2010; Nagelkerken *et al.*, 2015). This approach will have important consequences for how we rank the conservation of different marine habitats. In tropical seascapes, for example, mangroves, seagrasses and sandy shorelines are typically under-represented in marine conservation networks (relative to coral reefs) (Mumby, 2006; Unsworth & Cullen, 2010; Nagelkerken *et al.*, 2015). This integrates well with the gradual conceptual shift in perspective for marine conservation, from a historical focus on individual patches or habitats (Boström *et al.*, 2011) to the broader consideration of whole ecosystems transferring concepts and techniques from terrestrial landscape ecology.

A stronger focus on connectivity in conservation will likely benefit more than just fish populations and the performance of

marine reserves. It may, for example, also enhance the joint capacity of adjacent habitats, such as reefs, mangroves and seagrasses, to buffer seawater pH (e.g. Unsworth *et al.*, 2012), cope with sea-level rise (Saunders *et al.*, 2014), mitigate physical disturbance (e.g. Mumby & Hastings, 2008) and promote ecological resilience across seascapes (e.g. Olds *et al.*, 2012). At a broader scale, conserving connectivity will also enable organisms to access refugia from perturbations operating at local to regional scales (e.g. mining; Schlacher *et al.*, 2014) to global phenomena (e.g. rising temperatures; Magris *et al.*, 2014). Connectivity may also increase the value of ecosystem services provided by seascapes. For example, the value of a coral reef that is well connected to mangroves may be very different from that of a reef which is isolated from complementary resources (Hyndes *et al.*, 2014).

FUTURE DIRECTIONS

To date, all studies that have examined whether connectivity benefits conservation have been conducted in reef seascapes (eight on coral reefs; two on rocky reefs) (Appendix S2). It remains to be seen whether similar effects can also be demonstrated in a wider range of seascapes (e.g. coastal wetlands, soft sediments) and at the terrestrial–marine interface (i.e. beaches and estuaries) (e.g. Langlois *et al.*, 2005) (priority question 3, Table 1). Research has also largely focused on fishes and should now be expanded to evaluate the conservation benefits of connectivity for other organisms, such as crustaceans (e.g. MacArthur *et al.*, 2011) (priority question 4, Table 1).

Of the studies reviewed here, few had attempted to address questions about possible joint functional effects of seascape connectivity and marine reserves on ecological processes (e.g. herbivory, predation, recruitment). Consequently, we do not know whether effects of connectivity on ecological functions are common in reserve networks (e.g. Valentine *et al.*, 2008) (priority question 5, Table 1). This is a promising avenue for future research because the influence of connectivity on ecological processes provides the basis for conservation to influence ecosystem resistance and recovery (e.g. Olds *et al.*, 2012) (priority question 6, Table 1).

Studying the effect of seascape connectivity on animal movement will shed light on the mechanisms that underpin such spatial ecological relationships, and is key to appreciating how seascape conservation influences dispersal (e.g. Harrison *et al.*, 2012). At present, little is known about how the characteristics of seascapes at reserve boundaries (i.e. whether habitat is continuous or patchy) influence movement and dispersal from reserves (Freeman *et al.*, 2009; Pillans *et al.*, 2014; Pittman *et al.*, 2014). Spillover, which is a fisheries objective for many reserves (Halpern *et al.*, 2009), will likely be enhanced where habitat is continuous at reserve boundaries, but this will also work against the accumulation of fish biomass in reserves, which is a key conservation objective (Babcock *et al.*, 2012; Edgar *et al.*, 2014). Studies are needed to measure how the characteristics of seascapes at reserve boundaries influence the frequency and magnitude of spillover from reserves (e.g. Freeman *et al.*, 2009;

Pittman *et al.*, 2014) (priority question 7, Table 1). This will enhance our understanding of the capacity of reserves to achieve a common fisheries objective and thereby improve planning decisions about optimizing reserve design and placement. Given that reserves affect the behaviour and movement of exploited species (Halpern *et al.*, 2009; Babcock *et al.*, 2010), it will be important to examine whether effects of seascape connectivity on movement, dispersal and habitat use differ between seascapes within reserves, those that span reserve boundaries or areas that are open to fishing.

CONCLUSIONS

We find that the disciplines of landscape ecology and conservation biology lack close integration in the ocean, resulting in few studies that explicitly test for the effects of seascape connectivity on marine reserve performance. By contrast, landscape ecology has proven instrumental in terrestrial conservation (Rudnick *et al.*, 2012; Kool *et al.*, 2013), and we contend that it will offer similar benefits in the sea. Despite connectivity being increasingly viewed as important in marine conservation, this has generally not translated into quantitative objectives for conservation planning (Magris *et al.*, 2014). While this review focused on the implications of seascape connectivity for marine conservation, our findings may have broader implications for spatial conservation planning and ecosystem-based management because connectivity is rarely evaluated when assessing the effectiveness of management initiatives in either aquatic or terrestrial ecosystems. To better integrate connectivity into conservation we must focus research on identifying the mechanisms that underpin connectivity and the functions it delivers for ecosystems. Progress in spatial conservation planning will then be made by linking empirical studies of this nature with theoretical advances in our understanding of connectivity and the scale over which it impacts conservation outcomes.

ACKNOWLEDGEMENTS

This study was funded by the Australian Research Council, the Queensland Government and the Collaborative Research Network (CRN). S.J.P. was supported by NOAA's Coral Reef Conservation Program. We thank H. Faddy, B. Gilby and T. Martin for helpful discussions and comments on the manuscript. We are grateful to B. MacSharry and the United Nations Environment Programme World Conservation Monitoring Centre (UNEP-WCMC) for providing GIS layers for marine ecosystems and conservation areas.

REFERENCES

- Babcock, R.C., Shears, N.T., Alcala, A.C., Barrett, N.S., Edgar, G.J., Lafferty, K.D., McClanahan, T.R. & Russ, G.R. (2010) Decadal trends in marine reserves reveal differential rates of change in direct and indirect effects. *Proceedings of the National Academy of Sciences USA*, **107**, 18256–18261.

- Babcock, R.C., Egli, D. & Attwood, C.G. (2012) Individual-based model of reef fish populations in temperate marine reserves based on acoustic telemetry and census data. *Environmental Conservation*, **39**, 282–294.
- Beger, M., Linke, S., Watts, M., Game, E., Treml, E., Ball, I. & Possingham, H.P. (2010) Incorporating asymmetric connectivity into spatial decision making for conservation. *Conservation Letters*, **3**, 359–368.
- Bernhardt, J.R. & Leslie, H.M. (2013) Resilience to climate change in coastal marine ecosystems. *Annual Review of Marine Science*, **5**, 371–392.
- Boström, C., Pittman, S.J., Simenstad, C. & Kneib, R.T. (2011) Seascape ecology of coastal biogenic habitats: advances, gaps, and challenges. *Marine Ecology Progress Series*, **427**, 191–217.
- Cowen, R.K. & Sponaugle, S. (2009) Larval dispersal and marine population connectivity. *Annual Review of Marine Science*, **1**, 443–466.
- Edgar, G.J., Stuart-Smith, R.D., Willis, T.J. *et al.* (2014) Global conservation outcomes depend on marine protected areas with five key features. *Nature*, **506**, 216–220.
- Edwards, H.J., Elliott, I.A., Pressey, R.L. & Mumby, P.J. (2010) Incorporating ontogenetic dispersal, ecological processes and conservation zoning into reserve design. *Biological Conservation*, **143**, 457–470.
- Freeman, D.J., MacDiarmid, A.B. & Taylor, R.B. (2009) Habitat patches that cross marine reserve boundaries: consequences for the lobster *Jasus edwardsii*. *Marine Ecology Progress Series*, **388**, 159–167.
- Green, A.L., Maypa, A.P., Almany, G.R., Rhodes, K.L., Weeks, R., Abesamis, R.A., Gleason, M.G., Mumby, P.J. & White, A.T. (2014) Larval dispersal and movement patterns of coral reef fishes, and implications for marine reserve network design. *Biological Reviews*, doi:10.1111/brv.12155.
- Grober-Dunsmore, R., Pittman, S.J., Caldwell, C., Kendall, M.S. & Frazer, T.K. (2009) A landscape ecology approach for the study of ecological connectivity across tropical marine seascapes. *Ecological connectivity among tropical coastal ecosystems* (ed. by I. Nagelkerken), pp. 493–530. Springer, Heidelberg.
- Halpern, B., Lester, S. & Kellner, J. (2009) Spillover from marine reserves and the replenishment of fished stocks. *Environmental Conservation*, **36**, 268–276.
- Harrison, H.B., Williamson, D.H., Evans, R.D., Almany, G.R., Thorrold, S.R., Russ, G.R., Feldheim, K.A., Herwerden, L.V., Planes, S., Srinivasan, M., Berumen, M.L. & Jones, G.P. (2012) Larval export from marine reserves and the recruitment benefit for fish and fisheries. *Current Biology*, **22**, 1023–1028.
- Hodgson, J.A., Moilanen, A., Wintle, B.A. & Thomas, C.D. (2011) Habitat area, quality and connectivity: striking the balance for efficient conservation. *Journal of Applied Ecology*, **48**, 148–152.
- Huijbers, C.M., Connolly, R.M., Pitt, K.A., Schoeman, D.S., Schlacher, T.A., Burfeind, D.D., Steele, C., Olds, A.D., Maxwell, P.S., Babcock, R.C. & Rissik, D. (2014) Conservation benefits of marine reserves are undiminished near coastal rivers and cities. *Conservation Letters*, doi:10.1111/conl.12128.
- Huntington, B.E., Karnauskas, M., Babcock, E.A. & Lirman, D. (2010) Untangling natural seascape variation from marine reserve effects using a landscape approach. *PLoS ONE*, **5**, e12327.
- Hyndes, G.A., Nagelkerken, I., McLeod, R.J., Connolly, R.M., Lavery, P.S. & Vanderklift, M.A. (2014) Mechanisms and ecological role of carbon transfer within coastal seascapes. *Biological Reviews*, **89**, 232–254.
- Irlandi, E.A. & Crawford, M.K. (1997) Habitat linkages: the effect of intertidal saltmarshes and adjacent subtidal habitats on abundance, movement, and growth of an estuarine fish. *Oecologia*, **110**, 222–230.
- Jones, G.P. (2015) Mission impossible: unlocking the secrets of coral reef fish dispersal. *Ecology of fishes on coral reefs* (ed. by C. Mora), pp. 16–28. Cambridge University Press, Cambridge.
- Kool, J.T., Moilanen, A. & Treml, E.A. (2013) Population connectivity: recent advances and new perspectives. *Landscape Ecology*, **28**, 165–185.
- Langlois, T.J., Anderson, M.J., Babcock, R.C. & Kato, S. (2005) Reef-associated predators influence adjacent soft-sediment communities. *Ecology*, **86**, 1508–1519.
- Lester, S.E., Halpern, B.S., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B.I., Gaines, S.D., Aïramé, S. & Warner, R.R. (2009) Biological effects within no-take marine reserves: a global synthesis. *Marine Ecology Progress Series*, **384**, 33–46.
- MacArthur, L.D., Phillips, D.L., Hyndes, G.A., Hanson, C.E. & Vanderklift, M.A. (2011) Habitat surrounding patch reefs influences the diet and nutrition of the western rock lobster. *Marine Ecology Progress Series*, **436**, 191–205.
- MacNeil, M.A., Graham, N.A.J., Cinner, J.E., Wilson, S.K., Williams, I.D., Maina, J., Newman, S., Friedlander, A.M., Jupiter, S., Polunin, N.V.C. & McClanahan, T.R. (2015) Recovery potential of the world's coral reef fishes. *Nature*, **520**, 341–344.
- Magris, R.A., Pressey, R.L., Weeks, R. & Ban, N.C. (2014) Integrating connectivity and climate change into marine conservation planning. *Biological Conservation*, **170**, 207–221.
- Martin, T.S.H., Olds, A.D., Pitt, K.A., Johnston, A.B., Butler, I.R., Maxwell, P.S. & Connolly, R.M. (2015) Effective protection of fish on inshore coral reefs depends on the scale of mangrove-reef connectivity. *Marine Ecology Progress Series*, **527**, 157–165.
- Massol, F., Gravel, D., Mouquet, N., Cadotte, M.W., Fukami, T. & Leibold, M.A. (2011) Linking community and ecosystem dynamics through spatial ecology. *Ecology Letters*, **14**, 313–323.
- Micheli, F. & Peterson, C.H. (1999) Estuarine vegetated habitats as corridors for predator movements. *Conservation Biology*, **13**, 869–881.
- Moilanen, A., Wilson, K.A. & Possingham, H.P. (2009) *Spatial conservation prioritization: quantitative methods and computational tools*. Oxford University Press, Oxford.
- Mumby, P.J. (2006) Connectivity of reef fish between mangroves and coral reefs: algorithms for the design of marine reserves at seascape scales. *Biological Conservation*, **128**, 215–222.

- Mumby, P.J. & Hastings, A. (2008) The impact of ecosystem connectivity on coral reef resilience. *Journal of Applied Ecology*, **45**, 854–862.
- Nagelkerken, I., Grol, M.G.G. & Mumby, P.J. (2012) Effects of marine reserves versus nursery habitat availability on structure of reef fish communities. *PLoS ONE*, **7**, e36906.
- Nagelkerken, I., Sheaves, M., Baker, R. & Connolly, R.M. (2015) The seascape nursery: a novel spatial approach to identify and manage nurseries for coastal marine fauna. *Fish and Fisheries*, **16**, 362–371.
- Odum, E.P. (1968) Energy flow in ecosystems: a historical review. *American Zoologist*, **8**, 11–18.
- Olds, A.D., Pitt, K.A., Maxwell, P.S. & Connolly, R.M. (2012) Synergistic effects of reserves and connectivity on ecological resilience. *Journal of Applied Ecology*, **49**, 1195–1203.
- Olds, A.D., Albert, S., Maxwell, P.S., Pitt, K.A. & Connolly, R.M. (2013) Mangrove-reef connectivity promotes the functioning of marine reserves across the western Pacific. *Global Ecology and Biogeography*, **22**, 1040–1049.
- Olds, A.D., Connolly, R.M., Pitt, K.A., Maxwell, P.S., Aswani, S. & Albert, S. (2014) Incorporating surrogate species and seascape connectivity to improve marine conservation outcomes. *Conservation Biology*, **28**, 982–991.
- Pillans, R.D., Bearham, D., Boomer, A., Downie, R., Patterson, T.A., Thomson, D.P. & Babcock, R.C. (2014) Multi year observations reveal variability in residence of a tropical demersal fish, *Lethrinus nebulosus*: implications for spatial management. *PLoS ONE*, **9**, e105507.
- Pittman, S.J. & McAlpine, C.A. (2003) Movements of marine fish and decapod crustaceans: process, theory and application. *Advances in Marine Biology*, **44**, 205–294.
- Pittman, S.J. & Olds, A.D. (2015) *Seascape ecology of fishes on coral reefs*. Ecology of fishes on coral reefs (ed. by C. Mora), pp. 274–282. Cambridge University Press, Cambridge.
- Pittman, S.J., Monaco, M.E., Friedlander, A.M., Legare, B., Nemeth, R.S., Kendall, M.S., Poti, M., Clark, R.D., Wedding, L.M. & Caldwell, C. (2014) Fish with chips: tracking reef fish movements to evaluate size and connectivity of Caribbean marine protected areas. *PLoS ONE*, **9**, e96028.
- Rogers, A., Blanchard, J.L. & Mumby, P.J. (2014) Vulnerability of coral reef fisheries to a loss of structural complexity. *Current Biology*, **24**, 1000–1005.
- Rudnick, D.A., Ryan, S.J., Beier, P., Cushman, S.A., Dieffenbach, F., Epps, C.W., Gerber, L.R., Hartter, J., Jenness, J.S., Kintsch, J., Merelender, A.M., Perkl, R.M., Preziosi, D.V. & Trombulak, S.C. (2012) The role of landscape connectivity in planning and implementing conservation and restoration priorities. *Issues in Ecology*, **16**, 1–20.
- Salomon, A.K., Shears, N.T., Langlois, T.J. & Babcock, R. (2008) Cascading effects of fishing can alter carbon flow through a temperate coastal ecosystem. *Ecological Applications*, **18**, 1874–1887.
- Saunders, M.I., Leon, J.X., Callaghan, D.P., Roelfsema, C.M., Hamylton, S., Brown, C.J., Baldcock, T., Golshani, A., Phinn, S.R., Lovelock, C.E., Hoegh-Guldberg, O., Woodroffe, C.D. & Mumby, P.J. (2014) Interdependency of tropical marine ecosystems in response to climate change. *Nature Climate Change*, **4**, 724–729.
- Schlacher, T.A., Baco, A.R., Rowden, A.A., O'Hara, T.D., Clark, M.R., Kelley, C. & Dower, J.F. (2014) Seamount benthos in a cobalt-rich crust region of the central Pacific: conservation challenges for future seabed mining. *Diversity and Distributions*, **20**, 491–502.
- Sheaves, M. (2009) Consequences of ecological connectivity: the coastal ecosystem mosaic. *Marine Ecology Progress Series*, **391**, 107–115.
- Unsworth, R.K.F. & Cullen, L.C. (2010) Recognising the necessity for Indo-Pacific seagrass conservation. *Conservation Letters*, **3**, 63–73.
- Unsworth, R.K.F., Collier, C.J., Henderson, G.M. & McKenzie, L.J. (2012) Tropical seagrass meadows modify seawater carbon chemistry: implications for coral reefs impacted by ocean acidification. *Environmental Research Letters*, **7**, 024026.
- Valentine, J.F., Heck, K.L., Jr, Blackmon, D., Goecker, M.E., Christian, J., Kroutil, R.M., Peterson, B.J., Vanderklift, M.A., Kirsch, K.D. & Beck, M. (2008) Exploited species impacts on trophic linkages along reef–seagrass interfaces in the Florida keys. *Ecological Applications*, **18**, 1501–1515.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article at the publisher's web-site.

Appendix S1 Summary of seascape connectivity studies in reef and wetland systems.

Appendix S2 Summary of seascape connectivity studies that also quantified marine reserve effects, or were conducted inside marine reserves.

BIOSKETCH

Andrew Olds is a researcher at the University of the Sunshine Coast (Queensland, Australia). His research focuses on spatial ecology, conservation biology and the impacts of global change on marine ecosystems.

The research team includes ecologists from universities and research organizations in Australia, the United Kingdom, the United States of America and New Caledonia who are examining effects of seascape connectivity and marine reserves on faunal assemblages, ecological processes and ecosystem functioning. A.O., K.P. and R.C. conceived and designed the study. All authors contributed to synthesizing published data, analyses, interpreting the results and writing and revising the manuscript.

Editor: Derek Tittensor

APPENDIX: DATA SOURCES USED IN THIS STUDY

- Able, K.W., Balletto, J.H., Hagan, S.M., Jivoff, P.R. & Strait, K. (2007) Linkages between salt marshes and other nekton habitats in Delaware Bay, USA. *Reviews in Fisheries Science*, 15, 1–61.
- Able, K.W., Vivian, D.N., Petruzzelli, G. & Hagan, S.M. (2012a) Connectivity among salt marsh subhabitats: residency and movements of the mummichog (*Fundulus heteroclitus*). *Estuaries and Coasts*, 35, 743–753.
- Able, K.W., Grothues, T.M., Tumure, J.T., Byrne, D.M. & Clerkin, P. (2012b) Distribution, movements, and habitat use of small striped bass (*Morone saxatilis*) across multiple spatial scales. *Fishery Bulletin*, 110, 176–192.
- Aburto-Oropeza, O., Dominguez-Guerrero, I., Cota-Nieto, J. & Plomozo-Lugo, T. (2009) Recruitment and ontogenetic habitat shifts of the yellow snapper (*Lutjanus argentiventris*) in the Gulf of California. *Marine Biology*, 156, 2461–2472.
- Acosta, C.A. (1999) Benthic dispersal of Caribbean spiny lobsters among insular habitats: implications for the conservation of exploited marine species. *Conservation Biology*, 13, 603–612.
- Acosta, C.A. & Butler, M.J. (1997) Role of mangrove habitat as a nursery for juvenile spiny lobster, *Panulirus argus*. *Marine and Freshwater Research*, 48, 721–727.
- Aguiar-Perera, A. & Appeldoorn, R.S. (2007) Variation in juvenile fish density along the mangrove-seagrass-coral reef continuum in SW Puerto Rico. *Marine Ecology Progress Series*, 348, 139–148.
- Aller, E.A., Gullström, M., Eveleens Maarse, F.K.J., Gren, M., Nordlund, L.M., Jiddawi, N. & Eklöf, J.S. (2014) Single and joint effects of regional- and local-scale variables on tropical seagrass fish assemblages. *Marine Biology*, 161, 2395–2405.
- Baker, R., Fry, B., Rozas, L.P. & Minello, T.J. (2013) Hydrodynamic regulation of salt marsh contributions to aquatic food webs. *Marine Ecology Progress Series*, 490, 37–52.
- Barbier, E.B., Strand, I. & Sathirathai, S. (2002) Do open access conditions affect the valuation of an externality? Estimating the welfare effects of mangrove-fishery linkages in Thailand. *Environmental and Resource Economics*, 21, 343–367.
- Barbour, A.B., Adams, A.J. & Lorenzen, K. (2014) Size-based, seasonal, and multidirectional movements of an estuarine fish species in a habitat mosaic. *Marine Ecology Progress Series*, 507, 263–276.
- Bartholomew, A., Bohnsack, J.A., Smith, S.G., Ault, J.S., Harper, D.E. & McClellan, D.B. (2008) Influence of marine reserve size and boundary length on the initial response of exploited reef fishes in the Florida Keys National Marine Sanctuary, USA. *Landscape Ecology*, 23, 55–65.
- Beets, J., Muehlstein, L., Haught, K. & Schmitges, H. (2003) Habitat connectivity in coastal environments: patterns and movements of Caribbean coral reef fishes with emphasis on bluestriped grunt, *Haemulon sciurus*. *Gulf and Caribbean Research*, 14, 29–42.
- Beger, M. & Possingham, H. (2008) Environmental factors that influence the distribution of coral reef fishes: modeling occurrence data for broad-scale conservation and management. *Marine Ecology Progress Series*, 361, 1–13.
- Beger, M., Linke, S., Watts, M., Game, E., Treml, E., Ball, I. & Possingham, H.P. (2010) Incorporating asymmetric connectivity into spatial decision making for conservation. *Conservation Letters*, 3, 359–368.
- Berkström, C., Jørgensen, T.L. & Hellström, M. (2013a) Ecological connectivity and niche differentiation between two closely related fish species in the mangrove-seagrass-coral reef continuum. *Marine Ecology Progress Series*, 477, 201–215.
- Berkström, C., Lindberg, R., Thyresson, M. & Gullström, M. (2013b) Assessing connectivity in a tropical embayment: fish migrations and seascape ecology. *Biological Conservation*, 166, 43–53.
- Burkepile, D.E., Allgeier, J.E., Shantz, A.A., Pritchard, C.E., Lemoine, N.P., Bhatti, L.H. & Layman, C.A. (2013) Nutrient supply from fishes facilitates macroalgae and suppresses corals in a Caribbean coral reef ecosystem. *Scientific Reports*, 3, art. no. 1493.
- Campbell, S.J., Kartawijaya, T. & Sabarini, E.K. (2011) Connectivity in reef fish assemblages between seagrass and coral reef habitats. *Aquatic Biology*, 13, 65–77.
- Cerutti-Pereyra, F., Thums, M., Austin, C.M., Bradshaw, C.J.A., Stevens, J.R., Babcock, R.C., Pillans, R.D. & Meehan, M.G. (2014) Restricted movements of juvenile rays in the lagoon of Ningaloo Reef, Western Australia – evidence for the existence of a nursery. *Environmental Biology of Fishes*, 97, 371–383.
- Chateau, O. & Wantiez, L. (2007) Site fidelity and activity patterns of a humphead wrasse, *Cheilinus undulatus* (Labridae), as determined by acoustic telemetry. *Environmental Biology of Fishes*, 80, 503–508.
- Chin, A., Heupel, M.R., Simpfendorfer, C. & Tobin, A. (2013) Ontogenetic movements of juvenile blacktip reef sharks: evidence of dispersal and connectivity between coastal habitats and coral reefs. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 23, 468–474.
- Chittaro, P.M., Fryer, B.J. & Sale, P.F. (2004) Discrimination of French grunts (*Haemulon flavolineatum*, Desmarest, 1823) from mangrove and coral reef habitats using otolith microchemistry. *Journal of Experimental Marine Biology and Ecology*, 308, 169–183.
- Chittaro, P.M., Usseglio, P. & Sale, P.F. (2005) Variation in fish density, assemblage composition and relative rates of predation among mangrove, seagrass and coral reef habitats. *Environmental Biology of Fishes*, 72, 175–187.
- Claudet, J., García-Charton, J.A. & Lenfant, P. (2010) Combined effects of levels of protection and environmental variables at different spatial resolutions on fish assemblages in a marine protected area. *Conservation Biology*, 25, 105–114.
- Clynick, B. & Chapman, M.G. (2002) Assemblages of small fish in patchy mangrove forests in Sydney Harbour. *Marine and Freshwater Research*, 53, 669–677.
- Cocheret de la Morinière, E., Pollux, B.J.A., Nagelkerken, I. & Van der Velde, G. (2002) Post-settlement life cycle migration patterns and habitat preference of coral reef fish that use seagrass and mangrove habitats as nurseries. *Estuarine, Coastal and Shelf Science*, 55, 309–321.
- Davis, B., Johnston, R., Baker, R. & Sheaves, M. (2012) Fish utilisation of wetland nurseries with complex hydrological connectivity. *PLoS ONE*, 7, e49107. doi:10.1371/journal.pone.0049107.
- Davis, B., Baker, R. & Sheaves, M. (2014a) Seascape and metacommunity processes regulate fish assemblage structure in coastal wetlands. *Marine Ecology Progress Series*, 500, 187–202.
- Davis, B., Mattone, C. & Sheaves, M. (2014b) Bottom-up control regulates patterns of fish connectivity and assemblage structure in coastal wetlands. *Marine Ecology Progress Series*, 500, 175–186.
- Davis, J.P., Pitt, K.A., Fry, B., Olds, A.D. & Connolly, R.M. (2014c) Seascape-scale trophic links for fish on inshore coral reefs. *Coral Reefs*, 33, 897–907.
- Dennis, G.D., Smith-Vaniz, W.F., Colin, P.L., Hensley, D.A. & McGehee, M.A. (2005) Shore fishes from islands of the Mona Passage, Greater Antilles with comments on their zoogeography. *Caribbean Journal of Science*, 41, 716–743.
- Dorenbosch, M., van Riel, M.C., Nagelkerken, I. & Van Der Velde, G. (2004) The relationship of reef fish densities to the proximity of mangrove and seagrass nurseries. *Estuarine, Coastal and Shelf Science*, 60, 37–48.
- Dorenbosch, M., Grol, M., Nagelkerken, I. & Van der Velde, G. (2005a) Distribution of coral reef fishes along a coral reef-seagrass gradient: edge effects and habitat segregation. *Marine Ecology Progress Series*, 299, 277–288.
- Dorenbosch, M., Christianen, M.J.A., Nagelkerken, I. & Velde, G.V.D. (2005b) Indo-Pacific seagrass beds and mangroves contribute to fish density and diversity on adjacent coral reefs. *Marine Ecology Progress Series*, 302, 63–76.
- Dorenbosch, M., Grol, M.G.G., Nagelkerken, I. & Van der Velde, G. (2006a) Different surrounding landscapes may result in different fish assemblages in east African seagrass beds. *Hydrobiologia*, 563, 45–60.
- Dorenbosch, M., Grol, M.G.G., Nagelkerken, I. & Van der Velde, G. (2006b) Seagrass beds and mangroves as potential nurseries for the threatened Indo-Pacific humphead wrasse, *Cheilinus undulatus* and Caribbean rainbow parrotfish, *Scarus guacamaia*. *Biological Conservation*, 129, 277–282.
- Dorenbosch, M., Verberk, W., Nagelkerken, I. & van der Velde, G. (2007) Influence of habitat configuration on connectivity between fish assemblages of Caribbean seagrass beds, mangroves and coral reefs. *Marine Ecology Progress Series*, 334, 103–116.
- Dorenbosch, M., Grol, M.G.G., de Groene, A., van Der Velde, G. & Nagelkerken, I. (2009) Piscivore assemblages and predation pressure affect relative safety of some back-reef habitats for juvenile fish in a Caribbean bay. *Marine Ecology Progress Series*, 379, 181–196.
- Downie, R.A., Babcock, R.C., Thomson, D.P. & Vanderklift, M.A. (2013) Density of herbivorous fish and intensity of herbivory are influenced by proximity to coral reefs. *Marine Ecology Progress Series*, 482, 217–225.
- Dresser, B.K. & Kneib, R.T. (2007) Site fidelity and movement patterns of wild subadult red drum, *Sciaenops ocellatus* (Linnaeus), within a salt marsh-dominated estuarine landscape. *Fisheries Management and Ecology*, 14, 183–190.
- Drew, C.A. & Eggleston, D.B. (2008) Juvenile fish densities in Florida Keys mangroves correlate with landscape characteristics. *Marine Ecology Progress Series*, 362, 233–243.
- Eggleston, D.B., Dahlgren, C.P. & Johnson, E.G. (2004) Fish density, diversity, and size structure within multiple back reef habitats of Key West National Wildlife Refuge. *Bulletin of Marine Science*, 75, 175–204.
- Espino, F., González, J.A., Haroun, R. & Tuya, F. (2015) Abundance and biomass of the parrotfish *Sparisoma cretense* in seagrass meadows: temporal and spatial differences between seagrass interiors and seagrass adjacent to reefs. *Environmental Biology of Fishes*, 98, 121–133.
- Farmer, N.A. & Ault, J.S. (2011) Grouper and snapper movements and habitat use in Dry Tortugas, Florida. *Marine Ecology Progress Series*, 433, 169–184.
- Faunce, C.H. & Serafy, J.E. (2007) Nearshore habitat use by gray snapper (*Lutjanus griseus*) and bluestriped grunt (*Haemulon sciurus*): environmental gradients and ontogenetic shifts. *Bulletin of Marine Science*, 80, 473–495.
- Faunce, C.H. & Serafy, J.E. (2008) Selective use of mangrove shorelines by snappers, grunts, and great barracuda. *Marine Ecology Progress Series*, 356, 153–162.
- Fernandez, T.V., D'Anna, G., Badalamenti, F. & Pérez-Ruzafa, A. (2007) Habitat connectivity as a factor affecting fish assemblages in temperate reefs. *Aquatic Biology*, 1, 239–248.
- Fodrie, F.J. & Herzka, S.Z. (2008) Tracking juvenile fish movement and nursery contribution within arid coastal embayments via otolith microchemistry. *Marine Ecology Progress Series*, 361, 253–265.
- Fodrie, F.J. & Levin, L.A. (2008) Linking juvenile habitat utilization to population dynamics of California halibut. *Limnology and Oceanography*, 53, 799–812.
- Forcada, A., Bayle-Sempere, J.T., Valle, C. & Sánchez-Jerez, P. (2008) Habitat continuity effects on gradients of fish biomass across marine protected area boundaries. *Marine Environmental Research*, 66, 536–547.
- França, S., Costa, M.J. & Cabral, H.N. (2011) Inter- and intra-estuarine fish assemblage variability patterns along the Portuguese coast. *Estuarine, Coastal and Shelf Science*, 91, 262–271.
- Friedlander, A., Brown, E.K. & Monaco, M.E. (2007) Coupling ecology and GIS to evaluate efficacy of marine protected areas in Hawaii. *Ecological Applications*, 17, 715–730.
- García-Charton, J., Pérez-Ruzafa, A., Sánchez-Jerez, P., Bayle-Sempere, J., Renones, O. & Moreno, D. (2004) Multi-scale spatial heterogeneity, habitat structure, and the effect of marine reserves on western Mediterranean rocky reef fish assemblages. *Marine Biology*, 144, 161–182.
- García-Charton, J.A. & Pérez-Ruzafa, A. (1999) Ecological heterogeneity and evaluation of the effects of marine reserves. *Fisheries Research*, 42, 1–20.
- García-Charton, J.A. & Pérez-Ruzafa, A. (2001) Spatial pattern and the habitat structure of a Mediterranean rocky reef fish local assemblage. *Marine Biology*, 138, 917–934.
- García-Charton, J.A., Williams, I.D., Pérez-Ruzafa, A., Milazzo, M., Chemello, R., Marcos, C., Kitsos, M.S., Koukouras, A. & Riggio, S. (2000) Evaluating the ecological effects of Mediterranean marine protected areas: habitat, scale and the natural variability of ecosystems. *Environmental Conservation*, 27, 159–178.
- Gillanders, B.M. (1997) Patterns of abundance and size structure in the blue groper, *Achoerodus viridis* (Pisces, Labridae): evidence of links between estuaries and coastal reefs. *Environmental Biology of Fishes*, 49, 153–173.
- Gillanders, B.M. (2002) Connectivity between juvenile and adult fish populations: do adults remain near their recruitment estuaries? *Marine Ecology Progress Series*, 240, 215–223.
- Grabowski, J.H., Hughes, A.R., Kimbro, D.L. & Dolan, M.A. (2005) How habitat setting influences restored oyster reef communities. *Ecology*, 86, 1926–1935.
- Green, B.C., Smith, D.J. & Underwood, G.J.C. (2012a) Habitat connectivity and spatial complexity differentially affect mangrove and salt marsh fish assemblages. *Marine Ecology Progress Series*, 466, 177–192.
- Green, B.C., Smith, D.J., Grey, J. & Underwood, G.J.C. (2012b) High site fidelity and low site connectivity in temperate salt marsh fish populations: a stable isotope approach. *Oecologia*, 168, 245–255.
- Grober-Dunsmore, R., Frazer, T.K., Lindberg, W.J. & Beets, J. (2007) Reef fish and habitat relationships in a Caribbean seascape: the importance of reef context. *Coral Reefs*, 26, 201–216.
- Grober-Dunsmore, R., Frazer, T.K., Beets, J.P., Lindberg, W.J., Zwick, P. & Funicelli, N.A. (2008) Influence of landscape structure on reef fish assemblages. *Landscape Ecology*, 23, 47–53.
- Grober-Dunsmore, R., Bonito, V. & Aalbersberg, B. (2009) Findings from acoustic tagging reveal community based MPA in Fiji affords reasonable protection to letrichids. *Proceedings of the Inaugural Fiji Islands Conservation Science Forum, Suva, Fiji*.

- Grol, M., Nagelkerken, I., Rypel, A.L. & Layman, C.A. (2011) Simple ecological trade-offs give rise to emergent cross-ecosystem distributions of a coral reef fish. *Oecologia*, 165, 79–88.
- Grol, M.G.G., Rypel, A.L. & Nagelkerken, I. (2014) Growth potential and predation risk drive ontogenetic shifts among nursery habitats in a coral reef fish. *Marine Ecology Progress Series*, 502, 229–244.
- Gullström, M., Bodin, M., Nilsson, P.G. & Öhman, M.C. (2008) Seagrass structural complexity and landscape configuration as determinants of tropical fish assemblage composition. *Marine Ecology Progress Series*, 363, 241–255.
- Gullström, M., Berkstrom, C., Ohman, M., Bodin, M. & Dahlberg, M. (2011) Scale-dependent patterns of variability of a grazing parrotfish (*Leptoscarus vaigiensis*) in a tropical seagrass-dominated seascape. *Marine Biology*, 158, 1483–1495.
- Halpern, B.S. (2004) Are mangroves a limiting resource for two coral reef fishes? *Marine Ecology Progress Series*, 272, 93–98.
- Hamer, P.A., Jenkins, G.P. & Gillanders, B.M. (2005) Chemical tags in otoliths indicate the importance of local and distant settlement areas to populations of a temperate spard, *Pagrus auratus*. *Canadian Journal of Fisheries and Aquatic Sciences*, 62, 623–630.
- Hammerschlag, N. & Serafy, J.E. (2010) Nocturnal fish utilization of a subtropical mangrove–seagrass ecotone. *Marine Ecology*, 31, 364–374.
- Hammerschlag, N., Morgan, A.B. & Serafy, J.E. (2010a) Relative predation risk for fishes along a subtropical mangrove–seagrass ecotone. *Marine Ecology Progress Series*, 401, 259–267.
- Hammerschlag, N., Ovando, D. & Serafy, J.E. (2010b) Seasonal diet and feeding habits of juvenile fishes foraging along a subtropical marine ecotone. *Aquatic Biology*, 9, 279–290.
- Hammerschlag-Peyer, C.M. & Layman, C.A. (2012) Factors affecting resource use variation for an abundant coastal fish predator, *Lutjanus apodus*, in a Bahamian wetland system. *Bulletin of Marine Science*, 88, 211–230.
- Heithaus, E.R., Heithaus, P.A., Heithaus, M.R., Burkholder, D. & Layman, C.A. (2011) Trophic dynamics in a relatively pristine subtropical fringing mangrove community. *Marine Ecology Progress Series*, 428, 49–61.
- Hill, R.L., Appeldoorn, R.S., Murphy, B.R., Recksiek, C.W. & Prada, M. (2003) Scaling our understanding of habitat selection by coral reef fishery species. Proceedings of the Fifty-Fourth Annual Gulf and Caribbean Fisheries Institute, Providenciales, Turks and Caicos Islands (ed. by R.L. Creswell), pp. 566–579. GCFI, Marathon, FL.
- Hitt, S., Pittman, S.J. & Nemeth, R.S. (2011a) Diel movements of fishes linked to benthic seascape structure in a Caribbean coral reef ecosystem. *Marine Ecology Progress Series*, 427, 275–291.
- Hitt, S., Pittman, S.J. & Brown, K.A. (2011b) Tracking and mapping sun-synchronous migrations and diel space use patterns of *Haemulon sciurus* and *Lutjanus apodus* in the U.S. Virgin Islands. *Environmental Biology of Fishes*, 92, 525–538.
- Honda, K., Nakamura, Y., Nakaoka, M., Uy, W.H. & Fortes, M.D. (2013) Habitat use by fishes in coral reefs, seagrass beds and mangrove habitats in the Philippines. *PLoS ONE*, 8, e65735.
- Huijbers, C.M., Nagelkerken, I., Debrot, A.O. & Jongejans, E. (2013) Geographic coupling of juvenile and adult habitat shapes spatial population dynamics of a coral reef fish. *Ecology*, 94, 1859–1870.
- Huntington, B.E., Karnauskas, M., Babcock, E.A. & Lirman, D. (2010) Untangling natural seascape variation from marine reserve effects using a landscape approach. *PLoS ONE*, 5, e12327.
- Huxham, M., Kimani, E., Newton, J. & Augley, J. (2007) Stable isotope records from otoliths as tracers of fish migration in a mangrove system. *Journal of Fish Biology*, 70, 1554–1567.
- Irlandi, E.A. & Crawford, M.K. (1997) Habitat linkages: the effect of intertidal saltmarshes and adjacent subtidal habitats on abundance, movement, and growth of an estuarine fish. *Oecologia*, 110, 222–230.
- Jadot, C., Donnay, A., Acolas, M.L., Cornet, Y. & Begout Anras, M.L. (2006) Activity patterns, home-range size, and habitat utilization of *Sarpa salpa* (Teleostei: Sparidae) in the Mediterranean Sea. *ICES Journal of Marine Science*, 63, 128–139.
- Jelbart, J.E., Ross, P.M. & Connolly, R.M. (2007a) Fish assemblages in seagrass beds are influenced by the proximity of mangrove forests. *Marine Biology*, 150, 993–1002.
- Jelbart, J.E., Ross, P.M. & Connolly, R.M. (2007b) Patterns of small fish distributions in seagrass beds in a temperate Australian estuary. *Journal of the Marine Biological Association of the UK*, 87, 1297–1307.
- Johnston, R. & Sheaves, M. (2007) Small fish and crustaceans demonstrate a preference for particular small-scale habitats when mangrove forests are not accessible. *Journal of Experimental Marine Biology and Ecology*, 353, 164–179.
- Jones, D.L., Walter, J.F., Brooks, E.N. & Serafy, J.E. (2010) Connectivity through ontogeny: fish population linkages among mangrove and coral reef habitats. *Marine Ecology Progress Series*, 401, 245–258.
- Jones, K.M.M., McGrath, P.E. & Able, K.W. (2014) White perch *Morone americana* (Gmelin, 1789) habitat choice and movements: comparisons between *Phragmites*-invaded and *Spartina* reference marsh creeks based on acoustic telemetry. *Journal of Experimental Marine Biology and Ecology*, 455, 14–21.
- Kang, S.R. & King, S.L. (2013) Effects of hydrologic connectivity and environmental variables on nekton assemblage in a coastal marsh system. *Wetlands*, 33, 321–334.
- Kanouse, S., LaPeyre, M.K. & Nyman, J.A. (2007) Nekton use of *Ruppia maritima* and non-vegetated bottom habitat types within brackish marsh ponds. *Marine Ecology Progress Series*, 327, 61–69.
- Karnauskas, M., Huntington, B.E., Babcock, E.A. & Lirman, D. (2011) Pre-existing spatial patterns in fish abundances influence species-specific responses in a Caribbean marine reserve. *Marine Ecology Progress Series*, 432, 235–246.
- Karnauskas, M., Chérubin, L.M., Huntington, B.E., Babcock, E.A. & Thoney, D.A. (2012) Physical forces influence the trophic structure of reef fish communities on a remote atoll. *Limnology and Oceanography*, 57, 1403–1414.
- Kendall, M. (2005) A method for investigating seascape ecology of reef fish. *Proceedings of the Gulf and Caribbean Fisheries Institute*, 56, 356–366.
- Kendall, M., Christensen, J. & Hillis-Starr, Z. (2003) Multi-scale data used to analyze the spatial distribution of French grunts, *Haemulon flavolineatum*, relative to hard and soft bottom in a benthic landscape. *Environmental Biology of Fishes*, 66, 19–26.
- Kendall, M.S., Miller, T.J. & Pittman, S.J. (2011) Patterns of scale-dependency and the influence of map resolution on the seascape ecology of reef fish. *Marine Ecology Progress Series*, 427, 259–274.
- Kimirei, I.A., Nagelkerken, I., Griffioen, B., Wagner, C. & Mgaya, Y.D. (2011) Ontogenetic habitat use by mangrove/seagrass-associated coral reef fishes shows flexibility in time and space. *Estuarine, Coastal and Shelf Science*, 92, 47–58.
- Kimirei, I.A., Nagelkerken, I., Mgaya, Y.D. & Huijbers, C.M. (2013a) The mangrove nursery paradigm revisited: otolith stable isotopes support nursery-to-reef movements by Indo-Pacific fishes. *PLoS ONE*, 8, e66320.
- Kimirei, I.A., Nagelkerken, I., Trommelen, M., Blankers, P., van Hoytema, N., Hoeijmakers, D., Huijbers, C.M., Mgaya, Y.D. & Rypel, A.L. (2013b) What drives ontogenetic niche shifts of fishes in coral reef ecosystems? *Ecosystems*, 16, 783–796.
- Kimirei, I.A., Nagelkerken, I., Slooter, N., Gonzalez, E.T., Huijbers, C.M., Mgaya, Y.D. & Rypel, A.L. (2015) Demography of fish populations reveals new challenges in appraising juvenile habitat values. *Marine Ecology Progress Series*, 518, 225–237.
- Knib, R.T. & Wagner, S.L. (1994) Nekton use of vegetated marsh habitats at different stages of tidal inundation. *Marine Ecology Progress Series*, 106, 227–238.
- Knudby, A., Brenning, A. & LeDrew, E. (2010a) New approaches to modelling fish–habitat relationships. *Ecological Modelling*, 221, 503–511.
- Knudby, A., LeDrew, E. & Brenning, A. (2010b) Predictive mapping of reef fish species richness, diversity and biomass in Zanzibar using IKONOS imagery and machine-learning techniques. *Remote Sensing of Environment*, 114, 1230–1241.
- Knudby, A., Roelfsema, C., Lyons, M., Phinn, S. & Jupitser, S. (2011) Mapping fish community variables by integrating field and satellite data. object-based image analysis and modeling in a traditional Fijian fisheries management area. *Remote Sensing*, 3, 460–483.
- Koecb, B., Alos, J., Caro, A., Neveu, R., Cechlrou, R., Saragoin, G. & Lenfant, P. (2013) Contrasting fish behavior in artificial seascapes with implications for resources conservation. *PLoS ONE*, 8, e69303.
- Koivisto, M. & Westerborn, M. (2012) Invertebrate communities associated with blue mussel beds in a patchy environment: a landscape ecology approach. *Marine Ecology Progress Series*, 471, 101–110.
- Kopp, D., Bouchon-Navaro, Y., Louis, M. & Bouchon, C. (2007) Diel differences in the seagrass fish assemblages of a Caribbean island in relation to adjacent habitat types. *Aquatic Botany*, 87, 31–37.
- Krumme, U., Saint-Paul, U. & Rosenthal, H. (2004) Tidal and diel changes in the structure of a nekton assemblage in small intertidal mangrove creeks in northern Brazil. *Aquatic Living Resources*, 17, 215–229.
- Kuffner, I., Brock, J., Grober-Dunsmore, R., Bonito, V.E., Hickey, T.D. & Wright, C.W. (2007) Relationships between reef fish communities and remotely sensed rugosity measurements in Biscayne National Park, Florida, USA. *Environmental Biology of Fishes*, 78, 71–82.
- Langlois, T.J., Anderson, M.J., Babcock, R.C. & Kato, S. (2005) Reef-associated predators influence adjacent soft-sediment communities. *Ecology*, 86, 1508–1519.
- Langlois, T.J., Anderson, M.J., Babcock, R.C. & Kato, S. (2006) Marine reserves demonstrate trophic interactions across habitats. *Oecologia*, 147, 134–140.
- Layman, C.A., Arrington, D.A., Langerhans, R.B. & Silliman, B.R. (2004) Degree of fragmentation affects fish assemblage structure in Andros Island (Bahamas) estuaries. *Caribbean Journal of Science*, 40, 232–244.
- Layman, C.A., Quattrochi, J.P., Peyer, C.M. & Allgeier, J.E. (2007) Niche width collapse in a resilient top predator following ecosystem fragmentation. *Ecology Letters*, 10, 937–944.
- Layman, C.A., Allgeier, J.E., Yeager, L.A. & Stoner, E.W. (2013) Thresholds of ecosystem response to nutrient enrichment from fish aggregations. *Ecology*, 94, 530–536.
- Lugendo, B.R., Pronker, A., Cornelissen, I., Groene, A.D., Nagelkerken, I., Dorenbosch, M., Velde, G.V.D. & Mgaya, Y.D. (2005) Habitat utilisation by juveniles of commercially important fish species in a marine embayment in Zanzibar, Tanzania. *Aquatic Living Resources*, 18, 149–158.
- Lugendo, B.R., Nagelkerken, I., Kruitwagen, G., van der Velde, G. & Mgaya, Y.D. (2007) Relative importance of mangroves as feeding habitats for fishes: a comparison between mangrove habitats with different settings. *Bulletin of Marine Science*, 80, 497–512.
- Luo, J., Serafy, J.E., Sponaugle, S., Teare, P.B. & Kieckbusch, D. (2009) Movement of gray snapper *Lutjanus griseus* among subtropical seagrass, mangrove, and coral reef habitats. *Marine Ecology Progress Series*, 380, 255–269.
- MacArthur, L., Babcock, R. & Hyndes, G. (2008) Movements of the western rock lobster (*Panulirus cygnus*) within shallow coastal waters using acoustic telemetry. *Marine and Freshwater Research*, 59, 603–613.
- MacArthur, L.D., Phillips, D.L., Hyndes, G.A., Hanson, C.E. & Vanderklift, M.A. (2011) Habitat surrounding patch reefs influences the diet and nutrition of the western rock lobster. *Marine Ecology Progress Series*, 436, 191–205.
- Machemer, E.G.P., Walter, J.F., Serafy, J.E. & Kerstetter, D.W. (2012) Importance of mangrove shorelines for rainbow parrotfish *Scarus guacamaia*: habitat suitability modeling in a subtropical bay. *Aquatic Ecology*, 15, 87–98.
- McMahon, K.W., Berumen, M.L., Mateo, I., Elsdon, T.S. & Thorrold, S.R. (2011) Carbon isotopes in otolith amino acids identify residency of juvenile snapper (family: Lutjanidae) in coastal nurseries. *Coral Reefs*, 30, 1135–1145.
- McMahon, K.W., Berumen, M.L. & Thorrold, S.R. (2012) Linking habitat mosaics and connectivity in a coral reef seascape. *Proceedings of the National Academy of Sciences USA*, 109, 15372–15376.
- Madin, E.M.P., Madin, J.S. & Booth, D.J. (2011) Landscape of fear visible from space. *Scientific Reports*, 1, art. no. 14. doi:10.1038/srep00014.
- Manson, F.J., Loneragan, N.R., Harch, B.D., Skilleter, G.A. & Williams, L. (2005) A broad-scale analysis of links between coastal fisheries production and mangrove extent: a case-study for northeastern Australia. *Fisheries Research*, 74, 69–85.
- Martin, T.S.H., Olds, A.D., Pitt, K.A., Johnston, A.B., Butler, I.R., Maxwell, P.S. & Connolly, R.M. (2015) Effective protection of fish on inshore coral reefs depends on the scale of mangrove-reef connectivity. *Marine Ecology Progress Series*, 527, 157–165.
- Martosubroto, P. & Naamin, N. (1977) Relationship between tidal forests (mangroves) and commercial shrimp production in Indonesia. *Marine Research in Indonesia*, 18, 81–86.
- Mateo, I., Durbin, E., Appeldoorn, R., Adams, A., Juanes, F., Kingsley, R., Swart, P. & Durant, D. (2010) Role of mangroves as nurseries for French grunt *Haemulon flavolineatum* and schoolmaster *Lutjanus apodus* assessed by otolith elemental fingerprints. *Marine Ecology Progress Series*, 402, 197–212.
- Mellin, C., Andrefoet, S. & Ponton, D. (2007) Spatial predictability of juvenile fish species richness and abundance in a coral reef environment. *Coral Reefs*, 26, 895–907.
- Mellin, C., Bradshaw, C.J.A., Meekan, M.G. & Caley, M.J. (2010a) Environmental and spatial predictors of species richness and abundance in coral reef fishes. *Global Ecology and Biogeography*, 19, 212–222.
- Mellin, C., Huchery, C., Caley, M.J., Meekan, M.G. & Bradshaw, C.J.A. (2010b) Reef size and isolation determine the temporal stability of coral reef fish populations. *Ecology*, 91, 3138–3145.
- Meyer, C.G., Papastamatiou, Y.P. & Clark, T.B. (2010) Differential movement patterns and site fidelity among trophic groups of reef fishes in a Hawaiian marine protected area. *Marine Biology*, 157, 1499–1511.
- Meynecke, J., Lee, S. & Duke, N. (2008) Linking spatial metrics and fish catch reveals the importance of coastal wetland connectivity to inshore fisheries in Queensland, Australia. *Biological Conservation*, 141, 981–996.

- Micheli, F. & Peterson, C.H. (1999) Estuarine vegetated habitats as corridors for predator movements. *Conservation Biology*, 13, 869–881.
- Miller, M.W. & Gerstner, C.L. (2002) Reefs of an uninhabited Caribbean island: fishes, benthic habitat, and opportunities to discern reef fishery impact. *Biological Conservation*, 106, 37–44.
- Moore, C.H., Harvey, E.S. & Niel, K. (2010) The application of predicted habitat models to investigate the spatial ecology of demersal fish assemblages. *Marine Biology*, 157, 2717–2729.
- Mumby, P., Edwards, A., Arias-Gonzalez, J., Lindeman, K., Blackwell, P., Gall, A., Gorzczynska, M., Harborne, A., Perscod, C., Renken, H., Wabnitz, C. & Llewellyn, G. (2004) Mangroves enhance the biomass of coral reef fish communities in the Caribbean. *Nature*, 427, 533–536.
- Mumby, P.J. & Hastings, A. (2008) The impact of ecosystem connectivity on coral reef resilience. *Journal of Applied Ecology*, 45, 854–862.
- Nagelkerken, I. & Faunce, C.H. (2007) Colonisation of artificial mangroves by reef fishes in a marine seascape. *Estuarine, Coastal and Shelf Science*, 75, 417–422.
- Nagelkerken, I. & Van der Velde, G. (2004) Relative importance of interlinked mangroves and seagrass beds as feeding habitats for juvenile reef fish on a Caribbean island. *Marine Ecology Progress Series*, 274, 153–159.
- Nagelkerken, I., Dorenbosch, M., Verberk, W., Cocheret de la Moriniere, E. & van der Velde, G. (2000a) Day-night shifts of fishes between shallow-water biotopes of a Caribbean bay, with emphasis on the nocturnal feeding of Haemulidae and Lutjanidae. *Marine Ecology Progress Series*, 194, 55–64.
- Nagelkerken, I., Dorenbosch, M., Verberk, W., de la Moriniere, E. & van der Velde, G. (2000b) Importance of shallow-water biotopes of a Caribbean bay for juvenile coral reef fishes: patterns in biotope association, community structure and spatial distribution. *Marine Ecology Progress Series*, 202, 175–192.
- Nagelkerken, I., Kleijnen, S., Klop, T., van den Brand, R.A.C.J., Cocheret de la Moriniere, E. & van der Velde, G. (2001) Dependence of Caribbean reef fishes on mangroves and seagrass beds as nursery habitats: a comparison of fish faunas between bays with and without mangroves/seagrass beds. *Marine Ecology Progress Series*, 214, 225–235.
- Nagelkerken, I., Roberts, C.M., van der Velde, G., Dorenbosch, M., van Riel, M.C., Cocheret de la Moriniere, E. & Nienhuis, P.H. (2002) How important are mangroves and seagrass beds for coral-reef fish? The nursery hypothesis tested on an island scale. *Marine Ecology Progress Series*, 244, 299–305.
- Nagelkerken, I., Bothwell, J., Nemeth, R., Pitt, J. & van der Velde, G. (2008) Interlinkage between Caribbean coral reefs and seagrass beds through feeding migrations by grunts (Haemulidae) depends on habitat accessibility. *Marine Ecology Progress Series*, 368, 155–164.
- Nagelkerken, I., Grol, M.G.G. & Mumby, P.J. (2012) Effects of marine reserves versus nursery habitat availability on structure of reef fish communities. *PLoS ONE*, 7, e36906.
- Nakamura, Y. & Sano, M. (2004) Overlaps in habitat use of fishes between a seagrass bed and adjacent coral and sand areas at Amitoro Bay, Iriomote Island, Japan: importance of the seagrass bed as juvenile habitat. *Fisheries Science*, 70, 788–803.
- Nakamura, Y., Horinouchi, M., Shibuno, T., Tanaka, Y., Miyajima, T., Koike, I., Kurokura, H. & Sano, M. (2008) Evidence of ontogenetic migration from mangroves to coral reefs by black-tail snapper *Lutjanus fulvus*: stable isotope approach. *Marine Ecology Progress Series*, 355, 257–266.
- Nakamura, Y., Shibuno, T., Lecchini, D. & Watanabe, Y. (2009) Habitat selection by emperor fish larvae. *Marine Ecology Progress Series*, 6, 61–65.
- Nickerson, D.J. (1999) Trade-offs of mangrove area development in the Philippines. *Ecological Economics*, 28, 279–298.
- Olds, A.D., Connolly, R.M., Pitt, K.A. & Maxwell, P.S. (2012a) Primacy of seascape connectivity effects in structuring reef fish assemblages. *Marine Ecology Progress Series*, 462, 191–203.
- Olds, A.D., Connolly, R.M., Pitt, K.A. & Maxwell, P.S. (2012b) Habitat connectivity improves reserve performance. *Conservation Letters*, 5, 56–63.
- Olds, A.D., Pitt, K.A., Maxwell, P.S. & Connolly, R.M. (2012c) Synergistic effects of reserves and connectivity on ecological resilience. *Journal of Applied Ecology*, 49, 1195–1203.
- Olds, A.D., Albert, S., Maxwell, P.S., Pitt, K.A. & Connolly, R.M. (2013) Mangrove-reef connectivity promotes the functioning of marine reserves across the western Pacific. *Global Ecology and Biogeography*, 22, 1040–1049.
- Olds, A.D., Connolly, R.M., Pitt, K.A., Maxwell, P.S., Aswani, S. & Albert, S. (2014) Incorporating surrogate species and seascape connectivity to improve marine conservation outcomes. *Conservation Biology*, 28, 982–991.
- Pages, J.F., Bartumeus, F., Hereu, B., Lopez-Sanz, A., Romero, J. & Alcoverro, T. (2013) Evaluating a key herbivorous fish as a mobile link: a Brownian bridge approach. *Marine Ecology Progress Series*, 492, 199–210.
- Paw, J.N. & Chua, T.E. (1991) An assessment of the ecological and economic impact of mangrove conversion in Southeast Asia. *Towards an integrated management of tropical coastal resources. ICLARM Conference Proceedings*, 22, 201–212.
- Payne, N.L. & Gillanders, B.M. (2009) Assemblages of fish along a mangrove-mudflat gradient in temperate Australia. *Marine and Freshwater Research*, 60, 1–13.
- Payne, N.L., Snelling, E.P., Semmens, J.M. & Gillanders, B.M. (2013) Mechanisms of population structuring in giant Australian cuttlefish *Sepia apama*. *PLoS ONE*, 8, e58694.
- Penrose, H.M. (2011) Arid zone estuaries: nekton and trophic connectivity over heterogeneous landscapes. University of Queensland, Brisbane.
- Peterson, B.J., Valentine, J.F. & Heck, K.L., Jr (2013) The snapper–grunt pump: habitat modification and facilitation of the associated benthic plant communities by reef-resident fish. *Journal of Experimental Marine Biology and Ecology*, 441, 50–54.
- Pillars, R.D., Bearham, D., Boomer, A., Downie, R., Patterson, T.A., Thomson, D.P. & Babcock, R.C. (2014) Multi year observations reveal variability in residence of a tropical demersal fish, *Lethrinus nebulosus*: implications for spatial management. *PLoS ONE*, 9, e105507.
- Pittman, S., Caldwell, C., Hile, S. & Monaco, M. (2007a) Using seascape types to explain the spatial patterns of fish in the mangroves of SW Puerto Rico. *Marine Ecology Progress Series*, 348, 273–284.
- Pittman, S.J. & Brown, K.A. (2011) Multi-scale approach for predicting fish species distributions across coral reef seascapes. *PLoS ONE*, 6, e20583.
- Pittman, S.J., McAlpine, C.A. & Pittman, K.M. (2004) Linking fish and prawns to their environment: a hierarchical landscape approach. *Marine Ecology Progress Series*, 283, 233–254.
- Pittman, S.J., Christensen, J.D., Caldwell, C., Menza, C. & Monaco, M.E. (2007b) Predictive mapping of fish species richness across shallow-water seascapes in the Caribbean. *Ecological Modelling*, 204, 9–21.
- Pittman, S.J., Costa, B.M. & Battista, T.A. (2009) Using lidar bathymetry and boosted regression trees to predict the diversity and abundance of fish and corals. *Journal of Coastal Research*, 53, 27–38.
- Pittman, S.J., Monaco, M.E., Friedlander, A.M., Legare, B., Nemeth, R.S., Kendall, M.S., Poti, M., Clark, R.D., Wedding, L.M. & Caldwell, C. (2014) Fish with chips: tracking reef fish movements to evaluate size and connectivity of Caribbean marine protected areas. *PLoS ONE*, 9, e96028.
- Purkis, S.J., Graham, N.A.J. & Riegl, B.M. (2008) Predictability of reef fish diversity and abundance using remote sensing data in Diego Garcia (Chagos Archipelago). *Coral Reefs*, 27, 167–178.
- Randall, J.E. (1965) Grazing effect on sea grasses by herbivorous reef fishes in the West Indies. *Ecology*, 46, 255–260.
- Reid Nelson, T., Sutton, D. & DeVries, D.R. (2014) Summer movements of the gulf killifish (*Fundulus grandis*) in a northern Gulf of Mexico salt marsh. *Estuaries and Coasts*, 37, 1295–1300.
- Reis-Santos, P., Gillanders, B.M., Tanner, S.E., Vasconcelos, R.P., Elsdon, T.S. & Cabral, H.N. (2012) Temporal variability in estuarine fish otolith elemental fingerprints: implications for connectivity assessments. *Estuarine, Coastal and Shelf Science*, 112, 216–224.
- Reis-Santos, P., Tanner, S.E., Vasconcelos, R.P., Elsdon, T.S., Cabral, H.N. & Gillanders, B.M. (2013) Connectivity between estuarine and coastal fish populations: contributions of estuaries are not consistent over time. *Marine Ecology Progress Series*, 491, 177–183.
- Rozas, L.P. & Minello, T.J. (2006) Nekton use of *Vallisneria americana* Michx. (wild celery) beds and adjacent habitats in coastal Louisiana. *Estuaries and Coasts*, 29, 297–310.
- Rozas, L.P. & Odum, W.E. (1987) The role of submerged aquatic vegetation in influencing the abundance of nekton on contiguous tidal freshwater marshes. *Journal of Experimental Marine Biology and Ecology*, 114, 289–300.
- Rozas, L.P., Minello, T.J. & Dantin, D.D. (2012) Use of shallow lagoon habitats by nekton of the northeastern Gulf of Mexico. *Wetlands*, 35, 572–586.
- Rozas, L.P., Martin, C.W. & Valentine, J.F. (2013) Effects of reduced hydrological connectivity on the nursery use of shallow estuarine habitats within a river delta. *Marine Ecology Progress Series*, 492, 9–20.
- Russell, D. & McDougall, A. (2005) Movement and juvenile recruitment of mangrove jack, *Lutjanus argentimaculatus* (Forsskal), in northern Australia. *Marine and Freshwater Research*, 56, 465–475.
- Rypel, A.L. & Layman, C.A. (2008) Degree of aquatic ecosystem fragmentation predicts population characteristics of gray snapper (*Lutjanus griseus*) in Caribbean tidal creeks. *Canadian Journal of Fisheries and Aquatic Sciences*, 65, 335–339.
- Schultz, A.L., Malcolm, H.A., Bucher, D.J. & Smith, S.D.A. (2012) Effects of reef proximity on the structure of fish assemblages of unconsolidated substrata. *PLoS ONE*, 7, e49437.
- Sheaves, M. & Johnston, R. (2008) Influence of marine and freshwater connectivity on the dynamics of subtropical estuarine wetland fish metapopulations. *Marine Ecology Progress Series*, 357, 225–243.
- Sheaves, M. & Johnston, R. (2009) Ecological drivers of spatial variability among fish fauna of 21 tropical Australian estuaries. *Marine Ecology Progress Series*, 385, 245–260.
- Sheaves, M., Abrantes, K. & Johnston, R. (2007) Nursery ground value of an endangered wetland to juvenile shrimps. *Wetlands Ecol Manage*, 15, 311–327.
- Sheaves, M., Johnston, R., Connolly, R.M. & Baker, R. (2012) Importance of estuarine mangroves to juvenile banana prawns. *Estuarine, Coastal and Shelf Science*, 114, 208–219.
- Shibuno, T., Nakamura, Y., Horinouchi, M. & Sano, M. (2008) Habitat use patterns of fishes across the mangrove–seagrass–coral reef seascape at Ishigaki Island, southern Japan. *Ichthyological Research*, 55, 218–237.
- Skilleter, G.A., Olds, A., Loneragan, N.R. & Zharikov, Y. (2005) The value of patches of intertidal seagrass to prawns depends on their proximity to mangroves. *Marine Biology*, 147, 353–365.
- Staples, D.J., Vance, D.J. & Heales, D.S. (1985) Habitat requirements of juvenile penaeid prawns and their relationship to offshore fisheries. Second Australian National Prawn Seminar, Kooralybn, Queensland (ed. by P.C. Rothlisberg), pp. 47–54. CSIRO, Cleveland, Australia.
- Teo, S.L.H. & Able, K.W. (2003) Habitat use and movement of the mummichog (*Fundulus heteroclitus*) in a restored salt marsh. *Estuaries*, 26, 720–730.
- Thollot, P., Kulbicki, M. & Harmelin-Vivien, M. (1999) Trophic analysis and trophodynamic functioning of mangrove fish fauna of New Caledonia. *Comptes Rendus de Académie des Sciences, Paris*, 322, 607–619.
- Tupper, M.H. (2007) Spillover of commercially valuable reef fishes from marine protected areas in Guam, Micronesia. *Fishery Bulletin*, 105, 527–537.
- Tuya, F., Vanderklift, M.A., Hyndes, G.A., Wernberg, T., Thomsen, M.S. & Hanson, C. (2010) Proximity to rocky reefs alters the balance between positive and negative effects on seagrass fauna. *Marine Ecology Progress Series*, 405, 175–186.
- Tuya, F., Vanderklift, M.A., Wernberg, T. & Thomsen, M.S. (2011) Gradients in the number of species at reef–seagrass ecotones explained by gradients in abundance. *PLoS ONE*, 6, e20190.
- Unsworth, R., Bell, J. & Smith, D. (2007a) Tidal fish connectivity of reef and sea grass habitats in the Indo-Pacific. *Journal of the Marine Biological Association of the UK*, 87, 1287–1296.
- Unsworth, R., Wylie, E., Smith, D. & Bell, J. (2007b) Diel trophic structuring of seagrass bed fish assemblages in the Wakatobi Marine National Park, Indonesia. *Estuarine, Coastal and Shelf Science*, 72, 81–88.
- Unsworth, R.K.F., De León, P.S., Garrard, S.L., Jompa, J., Smith, D.J. & Bell, J.J. (2008) High connectivity of Indo-Pacific seagrass fish assemblages with mangrove and coral reef habitats. *Marine Ecology Progress Series*, 353, 213–224.
- Unsworth, R.K.F., Garrard, S.L., De León, P.S., Cullen, L.C., Smith, D.J., Sloman, K.A. & Bell, J.J. (2009) Structuring of Indo-Pacific fish assemblages along the mangrove–seagrass continuum. *Aquatic Biology*, 5, 85–95.
- Valentine, J.F., Heck, K.L., Jr, Blackmon, D., Goecker, M.E., Christian, J., Kroutil, R.M., Kirsch, K.D., Peterson, B.J., Beck, M. & Vanderklift, M.A. (2007) Food web interactions along seagrass–coral reef boundaries: effects of piscivore reductions on cross-habitat energy exchange. *Marine Ecology Progress Series*, 333, 37–50.
- Valentine, J.F., Heck, K.L., Jr, Blackmon, D., Goecker, M.E., Christian, J., Kroutil, R.M., Peterson, B.J., Vanderklift, M.A., Kirsch, K.D. & Beck, M. (2008) Exploited species impacts on trophic linkages along reef–seagrass interfaces in the Florida keys. *Ecological Applications*, 18, 1501–1515.
- Valentine-Rose, L. & Layman, C.A. (2011) Response of fish assemblage structure and function following restoration of two small Bahamian tidal creeks. *Restoration Ecology*, 19, 205–215.
- Vanderklift, M.A., How, J., Wernberg, T., MacArthur, L.D., Heck, K.L., Jr & Valentine, J.F. (2007) Proximity to reef influences density of small predatory fishes, while type of seagrass influences intensity of their predation on crabs. *Marine Ecology Progress Series*, 340, 235–243.

- Vasconcelos, R., Reis-Santos, P., Maia, A., Fonseca, V., Franca, S., Wouters, N., Costa, M.J. & Cabral, H.N. (2010) Nursery use patterns of commercially important marine fish species in estuarine systems along the Portuguese coast. *Estuarine, Coastal and Shelf Science*, 86, 613–624.
- Vasconcelos, R.P., Reis-Santos, P., Costa, M.J. & Cabral, H.N. (2011) Connectivity between estuaries and marine environment: integrating metrics to assess estuarine nursery function. *Ecological Indicators*, 11, 1123–1133.
- Vergés, A., Vanderklift, M.A., Doropoulos, C. & Hyndes, G.A. (2011) Spatial patterns in herbivory on a coral reef are influenced by structural complexity but not by algal traits. *PLoS ONE*, 6, e17115.
- Verweij, M.C. & Nagelkerken, I. (2007) Short and long-term movement and site fidelity of juvenile Haemulidae in back-reef habitats of a Caribbean embayment. *Hydrobiologia*, 592, 257–270.
- Verweij, M., Nagelkerken, I. & Hol, K. (2007) Space use of *Lutjanus apodus* including movement between a putative nursery and a coral reef. *Bulletin of Marine Science*, 81, 127–138.
- Verweij, M., Nagelkerken, I., Hans, I. & Ruseler, S.M. (2008) Seagrass nurseries contribute to coral reef fish populations. *Limnology and Oceanography*, 53, 1540–1547.
- Walsh, C.T., Reinfelds, I.V., Ives, M.C., Gray, C.A., West, R.J. & van der Meulen, D.E. (2013) Environmental influences on the spatial ecology and spawning behaviour of an estuarine-resident fish, *Macquaria colonorum*. *Estuarine, Coastal and Shelf Science*, 118, 60–71.
- Webb, S.R. & Kneib, R.T. (2002) Abundance and distribution of juvenile white shrimp *Litopenaeus setiferus* within a tidal marsh landscape. *Marine Ecology Progress Series*, 232, 213–223.
- Wedding, L.M. & Friedlander, A.M. (2008) Determining the influence of seascape structure on coral reef fishes in Hawaii using a geospatial approach. *Marine Geodesy*, 31, 246–266.
- Whaley, S., BurdJr, J.J. & Robertson, B. (2007) Using estuarine landscape structure to model distribution patterns in nekton communities and in juveniles of fishery species. *Marine Ecology Progress Series*, 330, 83–99.
- Yanez-Arancibia, A., Chavez, G.S. & Sanchez-Gil, P. (1985) Ecology of control mechanisms of natural fish production in the coastal zone. Fish community ecology in estuaries and coastal lagoons: towards an ecosystem integration (ed. by A. Yanez-Arancibia), pp. 571–594. UNAM Press, Mexico City.
- Yeager, L.A., Layman, C.A. & Allgeier, J.E. (2011) Effects of habitat heterogeneity at multiple spatial scales on fish community assembly. *Oecologia*, 167, 157–168.
- Yeager, L.A., Acevedo, C.L. & Layman, C.A. (2012) Effects of seascape context on condition, abundance, and secondary production of a coral reef fish, *Haemulon plumieri*. *Marine Ecology Progress Series*, 462, 231–240.